

## Literatuur - Gezamenlijke besluitvorming door gebruik van KomPas

1. Elwyn G, Frosch D, Thomson R, et al. Shared decision making: a model for clinical practice. *J Gen Intern Med.* 2012;27(10):1361-7.
2. Elwyn G, Frosch DL, Kobrin S. Implementing shared decision-making: consider all the consequences. *Implement Sci.* 2016;11:114
3. Reynolds A. Patient-centered Care. *Radiol Technol.* 2009;81(2):133-47.
4. Hoffmann TC, Montori VM, del Mar C. The connection between evidence-based medicine and shared decision making. *JAMA.* 2014;312(13):1295-6.
5. Moore CL, Kaplan SL. A framework and resources for shared decision making: opportunities for improved physical therapy outcomes. *Phys Ther.* 2018;98(12):1022-36.
6. Hoffmann TC, Lewis J, Maher CG. Shared decision making should be an integral part of physiotherapy practice. *Physiotherapy.* 2020;107:43-9.
7. Dierckx K, Deveugele M, Roosen P, et al. Implementation of shared decision making in physical therapy: observed level of involvement and patient preference. *Phys Ther.* 2013;93(10):1321-30.
8. Elwyn G, Dehlendorf C, Epstein RM, et al. Shared decision making and motivational interviewing: achieving patient-centered care across the spectrum of health care problems. *Ann Fam Med.* 2014;12(3):270-5.
9. Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: What does it mean? (or it takes at least two to tango). *Soc Sci Med.* 1997;44(5):681-92.
10. Jones LE, Roberts LC, Little PS, et al. Shared decision-making in back pain consultations: an illusion or reality? *Eur Spine J.* 2014;23(Suppl 1):S13-9.
11. Stenner R, Swinkels A, Mitchell T, et al. Exercise prescription for patients with non-specific chronic low back pain: a qualitative exploration of decision making in physiotherapy practice. *Physiotherapy.* 2016;102(4):332-8.
12. Sinnige A, Teijink JAW, Spruijt S, et al. The impact of personalized outcomes forecasts on clinical reasoning of physical therapists in Intermittent claudication: a vignette study. Submitted.
13. Aboyans V, Ricco JB, Bartelink MLEL, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS). *Eur Heart J.* 2018;39(9):763-816.
14. Conte MS, Pomposelli FB, Clair DG, et al. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: management of asymptomatic disease and claudication. *J Vasc Surg.* 2015;61(3 Suppl):2S-41S.
15. Sharma AM, Hazarika S, Solanki J, Annex BH. Angiogenesis in peripheral artery disease: focus on growth factor therapy. In: Emile R. Mohler, ER III, Annex BH (Eds.) *Regenerative Medicine for Peripheral Artery Disease*, pp. 1-16. Elsevier, 2016.
16. Merry AHH, Teijink JAW, Jongert MWA, et al. KNGF-richtlijn Symptomatisch perifeer arterieel vaatlijden. KNGF, 2014, revisie 2017. Geraadpleegd op 19-10-2021, van: <https://www.kngf2.nl/kennisplatform/richtlijnen/symptomatisch-perifeer-arterieel-vaatlijden>.
17. Barry MJ, Edgman-Levitan S. Shared decision making — the pinnacle of patient-centered care. *N Engl J Med.* 2012;366(9):780-1.
18. Sinnige A, Kittelson AJ, Wees PJ van der, et al. Personalized outcomes forecasts of supervised exercise therapy in intermittent claudication: an application of neighbors-based prediction methods with routinely collected clinical data. Submitted.
19. Kittelson AJ, Hoogeboom TJ, Schenkman M, et al. Person-centered care and physical therapy: a 'people-like-me' approach. *Phys Ther.* 2020;100(1):99-106.
20. Meerhoff GA, Dulmen SA van, Maas MJM, et al. Development and evaluation of an implementation strategy for collecting data in a national registry and the use of patient-reported outcome measures in physical therapist practices: quality improvement study. *Phys Ther.* 2017;97(8):837-51.
21. Dörenkamp S, Mesters I, Teijink J, et al. Difficulties of using single-diseased guidelines to treat patients with multiple diseases. *Front Public Health.* 2015;3:67
22. Ornstein SM, Nietert PJ, Jenkins RG, et al. The prevalence of chronic diseases and multimorbidity in primary care practice: a PPRNet report. *J Am Board Fam Med.* 2013;26(5):518-24.
23. Dörenkamp S, Mesters EPE, Nijhuis-van der Sanden MWG, et al. How well do randomized controlled trials reflect standard care: a comparison between scientific research data and standard care data in patients with intermittent claudication undergoing supervised exercise therapy. *PLoS One.* 2016;11(6):e0157921.
24. Buuren S van. Curve matching: a data-driven technique to improve individual prediction of childhood growth. *Ann Nutr Metab.* 2014;65(3):227-33.

25. Kruidenier LM, Nicolaï SP, Willigendael EM, et al. Functional claudication distance: a reliable and valid measurement to assess functional limitation in patients with intermittent claudication. *BMC Cardiovasc Disord.* 2009;9(9).
26. Gardner A, Skinner J, Cantwell B, et al. Progressive vs single-stage treadmill tests for evaluation of claudication. *Med Sci Sports Exerc.* 1991;23(4):402-8.
27. Nordanstig J, Wann-Hansson C, Karlsson J, et al. Vascular Quality of Life Questionnaire-6 facilitates health-related quality of life assessment in peripheral arterial disease. *J Vasc Surg.* 2014;59(3):700-7.
28. Cheema A, Bagchi R. The effect of goal visualization on goal pursuit: implications for consumers and managers. *J Mark.* 2011;75(2):109-23.